

हमारा विश्वास... हर एक विद्यार्थी है स्वास

JEE  
MAIN  
JAN'19

**QUESTION WITH SOLUTION**  
DATE : 10-01-2019 \_ EVENING

**IIT  
NIT**  
XI, XII & XII Pass

**AIIMS  
NEET**  
XI, XII & XII Pass

**BOARDS  
NTSE  
OLYMPIADS**  
V to X Class

RESIDENTIAL  
COACHING PROGRAM  
**rona**  
Discipline-Bridge between dreams & Success

**20000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**4626**

(Under 50000 Rank)

JEE (Main)

**13953**

NEET / AIIMS NTSE / OLYMPIADS

**662**

(since 2016)

**1066**

(5th to 10th class)

Toll Free :  
1800-212-1799

**MOTION™**

Nurturing potential through education

H.O. : 394, Rajeev Gandhi Nagar, Kota  
www.motion.ac.in |✉: info@motion.ac.in

# [CHEMISTRY]

1. The 71<sup>st</sup> electron of an element X with an atomic number of 71 enters into the orbital:  
 (A) 5d (B) 4f (C) 6p (D) 6s

Sol. **B**

2. The ground state energy of hydrogen atom is -13.6eV. The energy of second excited state of He<sup>+</sup> ion in eV is:  
 (A) -54.4 (B) -27.2 (C) -6.04 (D) -3.4

Sol. **C**

$$(E)_n^{\text{th}} = (E_{\text{GND}})_{\text{H}^+} \cdot \frac{Z^2}{n^2}$$

$$E_{3^{\text{rd}}}(\text{He}^+) = (-13.6\text{eV}) \cdot \frac{2^2}{3^2} = -6.04 \text{ eV}$$

3. An ideal gas undergoes isothermal compression from 5m<sup>3</sup> to 1 m<sup>3</sup> against a constant external pressure of 4Nm<sup>-2</sup>. Heat released in this process is used to increase the temperature of 1 mole of Al. If molar heat capacity of Al is 24 J mol<sup>-1</sup> K<sup>-1</sup>, the temperature of Al increases of by:

- (A) 2k (B)  $\frac{2}{3}$ K (C)  $\frac{3}{2}$ K (D) 1K

Sol. **B**

Work done on isothermal irreversible for ideal gas

$$\begin{aligned} &= -P_{\text{ext}}(V_2 - V_1) \\ &= -4 \text{ N/m}^2 (1\text{m}^3 - 5\text{m}^3) \\ &= 16 \text{ Nm} \end{aligned}$$

Isothermal process for ideal gas

$$\Delta U = 0$$

$$\begin{aligned} q &= -w \\ &= -16 \text{ Nm} \\ &= -16 \text{ J} \end{aligned}$$






Heat used to increase temperature of Al

$$q = n C_m \Delta T$$

$$16 \text{ J} = 1 \times 24 \frac{\text{J}}{\text{mol K}} \times \Delta T$$


$$\Delta T = \frac{2}{3} \text{ K}$$

4. The major product obtained in the following reaction is:



Sol. **B**

5. Among the following reactions of hydrogen with halogens, the one that requires a catalyst is:




Sol. D

Because reaction of  $\text{H}_2$  and  $\text{I}_2$  is Reversible in nature .

6. In the reaction of oxalate with permanganate in acidic medium, the number of electrons involved in producing one molecule of  $\text{CO}_2$  is:




Sol. A



$10e^-$  transfer 10 molecule of  $\text{CO}_2$  So per molecule of  $\text{CO}_2$  transfer of  $e^-$  is '1'

7. Which of the following tests cannot be used for identifying amino acids?



Sol. A

8. The pair that contains two P-H bonds in each of the oxoacids is:




Sol. C

9. For an elementary chemical reaction,  $\text{A}_2 \xrightleftharpoons[k_{-1}]{k_1} 2\text{A}$ , the expression for  $\frac{d[\text{A}]}{dt}$  is:



Sol. A



$$\frac{d[\text{A}]}{dt} = 2k_1[\text{A}_2] - 2k_{-1}[\text{A}]^2$$

10. Elevation in the boiling point for 1 molal solution of glucose is 2 K. the depression in the freezing point for 2 molal solution of glucose in the same solvent is 2 K. The relation between  $K_b$  and  $k_f$  is:




Sol. A

$$\frac{\Delta T_b}{\Delta T_f} = \frac{i \cdot m \times k_b}{i \times m \times k_f}$$

$$\frac{2}{2} = \frac{1 \times 1 \times k_b}{1 \times 2 \times k_f}$$

$$k_b = 2K_f$$

11. What is the IUPAC name of the following compound?



- (A) 3-Bromo-3-methyl-1, 2-dimethylprop-1-ene  
 (B) 3-Bromo-1, 2-dimethylbut-1-ene  
 (C) 4-Bromo-3-methylpent-2-ene  
 (D) 2-Bromo-3-methylpent-3-ene





Sol. C

12. Sodium metal on dissolution in liquid ammonia gives a deep blue solution due to the formation of:

- (A) ammoniated electrons (B) sodium ion-ammonia complex  
(C) sodium-ammonia complex (D) sodamide


Sol. A

13. An aromatic compound 'A' having molecular formula  $C_7H_6O_2$  on treating with aqueous ammonia and heating forms compound 'B'. The compound 'B' on reaction with molecular bromine and potassium hydroxide provides compound 'C' having molecular formula  $C_6H_7N$ . The structure of 'A' is:



Sol. D

14. In the cell






the cell potential is 0.92 V when a  $10^{-6}$  molal HCl solution is used. The standard electrode potential of (AgCl/AgCl<sup>-</sup>) electrode is:

(Given,  $\frac{2.303RT}{F} = 0.06 \text{ V}$  at 298 K)

- (A) 0.76V (B) 0.40 V (C) 0.94 V (D) 0.20 V

Sol. D



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.06}{2} \log_{10} (H^+)^2 (Cl^-)^2$$

$$.925 = \left( E_{H_2/H^+}^0 + E_{AgCl/Ag,Cl^-}^0 \right) - \frac{0.06}{2} \log_{10} ((10^{-6})^2 (10^{-6})^2)$$

$$.92 = 0 + E_{AgCl/Ag,Cl^-}^0 - 0.031 \log_{10} (10^{-6})^4$$

$$E_{AgCl/Ag,Cl^-}^0 = .92 + .03 \times -24 = 0.2 \text{ V}$$

15. A compound of formula  $A_2B_3$  has the hcp lattice. Which atom forms the hcp lattice and what fraction of tetrahedral voids is occupied by the other atoms:

- (A) hcp lattice-B,  $\frac{2}{3}$  Tetrahedral voids-A (B) hcp lattice-A,  $\frac{2}{3}$  Tetrahedral voids-B

- (C) hcp lattice-A,  $\frac{1}{3}$  Tetrahedral voids-B (D) hcp lattice-B,  $\frac{1}{3}$  Tetrahedral voids-A

Sol. C

$A_2B_3$  has HCP lattice

If A form HCP, then  $\frac{3^{\text{th}}}{4}$  of THV must occupied by B to form  $A_2B_3$






If B form HCP, then  $\frac{1^{\text{th}}}{3}$  of THV must occupied by A to form  $A_2B_3$

16. The difference in the number of unpaired electrons of a metal ion in its high-spin and low-spin octahedral complexes is two. The metal ion is:

(A)  $Mn^{2+}$  (B)  $Fe^{2+}$  (C)  $Co^{2+}$  (D)  $Ni^{2+}$






Sol. C

17. The major product of the following reaction is:



Sol. A/B (NTA)

18. What will be the major product in the following mononitration reaction?



Sol. D

19. 5.1 g  $NH_4SH$  is introduced in 3.0 L evacuated flask at  $327^\circ C$ . 30% of the solid  $NH_4SH$  decomposed to  $NH_3$  and  $H_2S$  as gases. The  $K_p$  of the reaction at  $327^\circ C$  is ( $R = 0.082 \text{ L atm mol}^{-1} \text{ K}^{-1}$ , Molar mass of S =  $32 \text{ g mol}^{-1}$ , molar mass of N =  $14 \text{ g mol}^{-1}$ )

(A)  $1 \times 10^{-4} \text{ atm}^2$  (B)  $4.9 \times 10^{-3} \text{ atm}^2$  (C)  $0.242 \times 10^{-4} \text{ atm}^2$  (D)  $0.242 \text{ atm}^2$

Sol. D



$$n = \frac{5.1}{51} = .1 \text{ mole} \quad 0 \quad 0$$

$$.1(-1-\alpha) \quad .1\alpha \quad .1\alpha$$

$$\alpha = 30\% = .3$$

so number of moles at equilibrium

$$.1(1 - .3) \quad .1 \times .3 \quad .1 \times .3$$

$$= .07 \quad =.03 \quad =.03$$

Now use PV = nRT at equilibrium

$$P_{\text{total}} \times 3 \text{ lit} = (.03 + .03) \times .082 \times 600$$

$$P_{\text{total}} = .984 \text{ atm}$$





At equilibrium

$$P_{\text{NH}_3} = P_{\text{H}_2\text{S}} = \frac{P_{\text{total}}}{2} = .492$$

$$\text{So } K_p = P_{\text{NH}_3} \cdot P_{\text{H}_2\text{S}} = (.492) (.492)$$





$$K_p = .242 \text{ atm}^2$$

20. The reaction that is NOT involved in the ozone layer depletion mechanism in the stratosphere is:







Sol. D

21. Which is the most suitable reagent for the following transformation?







Sol. C

22. The electrolytes usually used in the electroplating of gold and silver, respectively, are :




Sol. A





23. The process with negative entropy change is :



Sol. B



24. The number of 2-centre-2-electron and 3-centre-2-electron bonds in  $\text{B}_2\text{H}_6$ , respectively, are :



Sol. B

25. A reaction of cobalt (III) chloride and ethylenediamine in a 1 : 2 mole ratio generates two isomeric products A (violet coloured) and B (green coloured). A can show optical activity, but, B is optically inactive. What type of isomers does A and B represent ?



Sol. C

26. Haemoglobin and gold sol are examples of :  
 (A) negatively charged sols  
 (B) positively and negatively charged sols, respectively  
 (C) positively charged sols  
 (D) negatively and positively charged sols, respectively

Sol. **B**  
 Haemoglobin  $\longrightarrow$  positive sol  
 Ag  $-$ sol  $\longrightarrow$  negative sol

27. The amount of sugar ( $C_{12}H_{22}O_{11}$ ) required to prepare 2 L of its 0.1 M aqueous solution is :  
 (A) 136.8g (B) 68.4g (C) 17.1g (D) 34.2g


Sol. **B**  

$$\text{Molarity} = \frac{(n)_{\text{solute}}}{V_{\text{solution}} \text{ (in lit)}}$$

$$0.1 = \frac{\text{wt}/342}{2}$$

$$\text{wt} (C_{12}H_{22}O_{11}) = 68.4 \text{ gram}$$

28. The major product of the following reaction is :



Sol. **C**


29. The correct match between item 'I' and item 'II' is :

Item 'I' Compound	Item 'II' reagent
(A) Lysine	(P) 1-naphthol
(B) Furfural	(Q) ninhydrin
(C) Benzyl alcohol	(R) $KMnO_4$
(D) Styrene	(S) Ceric ammonium nitrate

(A) A  $\rightarrow$  Q ; B  $\rightarrow$  P ; C  $\rightarrow$  R ; D  $\rightarrow$  S  
 (B) A  $\rightarrow$  Q ; B  $\rightarrow$  R ; C  $\rightarrow$  S ; D  $\rightarrow$  P  
 (C) A  $\rightarrow$  R ; B  $\rightarrow$  P ; C  $\rightarrow$  Q ; D  $\rightarrow$  S  
 (D) A  $\rightarrow$  Q ; B  $\rightarrow$  P ; C  $\rightarrow$  S ; D  $\rightarrow$  R

Sol. **D**

30. The major product of the following reaction is :



Sol. **A**