

# JEE MAIN

QUESTION PAPER  
WITH SOLUTION

**38000+**  
SELECTIONS SINCE 2007



## PHYSICS

**26<sup>th</sup> June 2022 | Shift - 2**

# MOTION<sup>®</sup>

JEE (Main+Advanced) | NEET | NTSE | Olympiads | Boards

# Umeed **Rank** Ki Ho Ya **Selection** Ki, JEET NISCHIT HAI!

MOST PROMISING RANKS  
PRODUCED BY MOTION FACULTIES

NATION'S BEST SELECTION  
PERCENTAGE (%) RATIO

NEET / AIIMS

AIR-1 TO 10  
25 TIMES

AIR-11 TO 25  
37 TIMES

AIR-26 TO 50  
43 TIMES

AIR-51 TO 100  
78 TIMES

JEE MAIN+ADVANCED

AIR-1 TO 10  
8 TIMES

AIR-11 TO 25  
6 TIMES

AIR-26 TO 50  
18 TIMES

AIR-51 TO 100  
30 TIMES

**MOTION**<sup>®</sup>  
JEE | NEET | NTSE | BOARDS | OLYMPIADS



**NITIN VIJAY (NV Sir)**  
Founder & CEO

STUDENT  
QUALIFIED  
IN NEET

2021 3276 / 3411  
= 93.12%

2020 2663 / 2843  
= 93.66%

2019 2041 / 2212  
= 92.27%

STUDENT  
QUALIFIED IN  
JEE ADVANCED

2021 1256 / 2994  
= 41.95%

2020 994 / 2538  
= 39.16%

2019 769 / 2105  
= 36.53%

STUDENT  
QUALIFIED  
IN JEE MAIN

2021 2994 / 4087  
= 73.25%

2020 2538 / 3554  
= 71.44%

2019 2288 / 3316  
= 68.99%

### SECTION - A

1. The dimension of mutual inductance is :

- (A)  $[ML^2T^{-2}A^{-1}]$       (B)  $[ML^2T^{-3}A^{-1}]$       (C)  $[ML^2T^{-2}A^{-2}]$       (D)  $[ML^2T^{-3}A^{-2}]$

Sol. (C)

$e_2$  : induced emf in secondary coil

$i_1$  : Current in primary coil


$M$  : Mutual inductance

$$e_2 = -M \frac{di_1}{dt}$$

$$M = -\frac{e_2}{\frac{di_1}{dt}}$$

$$[M] = \frac{[e_2]}{\left[\frac{di_1}{dt}\right]} = \frac{\left[\frac{W}{q}\right]}{\left[\frac{[A]}{[T]}\right]} = \frac{[ML^2T^{-2}]}{[AT^{-1}]} = [ML^2T^{-2}A^{-2}]$$


2. In the arrangement shown in the figure  $a_1, a_2, a_3$  and  $a_4$  are the acceleration of masses  $m_1, m_2, m_3$  and  $m_4$  respectively. Which of the following relation is true for this arrangement?



- (A)  $4a_1 + 2a_2 + a_3 + a_4 = 0$   
 (C)  $a_1 + 4a_2 + 3a_3 + 2a_4 = 0$

- (B)  $a_1 + 4a_2 + 3a_3 + a_4 = 0$   
 (D)  $2a_1 + 2a_2 + 3a_3 + a_4 = 0$





Using constraint

$$\sum \vec{T} \cdot \vec{a} = 0$$

$$-4Ta_1 - 2Ta_2 - Ta_3 - Ta_4 = 0$$

$$4a_1 + 2a_2 + a_3 + a_4 = 0$$

3. Arrange the four graphs in descending order of total work done; where  $W_1, W_2, W_3$  and  $W_4$  are the work done corresponding to figure a, b, c and d respectively.




Figure-a




Figure-b




Figure-c




Figure-d

- (A)  $W_3 > W_2 > W_1 > W_4$   
 (C)  $W_2 > W_3 > W_4 > W_1$

- (B)  $W_3 > W_2 > W_4 > W_1$   
 (D)  $W_2 > W_3 > W_1 > W_4$





Sol. (A)

Work done = area under F - x curve. Area below x-axis is negative & area above x-axis is positive.  
So  $W_3 > W_2 > W_1 > W_4$

4. A solid spherical ball is rolling on a frictionless horizontal plane surface about its axis of symmetry. The ratio of rotational kinetic energy of the ball to its total kinetic energy is

- (A)  $\frac{2}{5}$                       (B)  $\frac{2}{7}$                       (C)  $\frac{1}{5}$                       (D)  $\frac{7}{10}$

Sol. (B)

$$K_{\text{total}} = K_{\text{rotational}} + K_{\text{Translational}}$$

$$K_{\text{total}} = \frac{1}{2} I_{\text{cm}} \omega^2 + \frac{1}{2} m v_{\text{cm}}^2$$

$$v_{\text{cm}} = R\omega \text{ for pure rolling}$$

$$I_{\text{cm}} = \frac{2}{5} m R^2$$

$$K_{\text{Rot}} = \frac{1}{2} I_{\text{cm}} \omega^2 = \frac{1}{2} \times \frac{2}{5} m R^2 \times \frac{v_{\text{cm}}^2}{R^2} = \frac{1}{5} m v_{\text{cm}}^2$$

$$K_{\text{total}} = \frac{1}{5} m v_{\text{cm}}^2 + \frac{1}{2} m v_{\text{cm}}^2 = \frac{7}{10} m v_{\text{cm}}^2$$

$$\frac{K_{\text{Rot}}}{K_{\text{Total}}} = \frac{\frac{1}{5} m v_{\text{cm}}^2}{\frac{7}{10} m v_{\text{cm}}^2} = \frac{2}{7}$$

5. Given below are two statements: One is labelled as Assertion A and the other is labelled as Reason R.


**Assertion A :** If we move from poles to equator, the direction of acceleration due to gravity of earth always points towards the center of earth without any variation in its magnitude.

**Reason R :** At equator, the direction of acceleration due to the gravity is towards the center of earth.

In the light of above statements, choose the correct answer from the options given below:

- (A) Both A and R are true and R is the correct explanation of A.  
(B) Both A and R are true but R is NOT the correct explanation of A.  
(C) A is true but R is false  
(D) A is false but R is true

Sol. (D)



Effective acceleration due to gravity is the resultant of g &  $r\omega^2$  whose direction & magnitude depends upon θ. hence assertion is false.

When  $\theta = 0^\circ$  (at equator), effective acceleration is radially inward.

6. If ρ is the density and η is coefficient of viscosity of fluid which flows with a speed v in the pipe of diameter d, the correct formula for Reynolds number  $R_e$  is:

- (A)  $R_e = \frac{\eta d}{\rho v}$                       (B)  $R_e = \frac{\rho v}{\eta d}$                       (C)  $R_e = \frac{\rho v d}{\eta}$                       (D)  $R_e = \frac{\eta}{\rho v d}$



Sol. (C)

Reynold's number is given by  $\frac{\rho v d}{\eta}$

7. A flask contains argon and oxygen in the ratio of 3 : 2 in mass and the mixture is kept at 27°C. The ratio of their average kinetic energy per molecule respectively will be:

- (A) 3 : 2                      (B) 9 : 4                      (C) 2 : 3                      (D) 1 : 1


Sol. **Bonus**

(D)

Average K.E. / molecule =  $\frac{f}{2} kT$


$$\text{So, } \frac{K_{Ar}}{K_{O_2}} = \frac{\frac{3}{2}kT}{\frac{5}{2}kT} = \frac{3}{5}$$

8. The charge on capacitor of capacitance 15μF in the figure given below is:



- (A) 60μc                      (B) 130μc                      (C) 260μc                      (D) 585μc

Sol. (A)



$$\frac{1}{C_{eq}} = \frac{1}{10} + \frac{1}{15} + \frac{1}{20} = \frac{12+8+6}{120} = \frac{26}{120}$$

$$C_{eq} = \frac{60}{13} \mu F$$


$$Q = \frac{13 \times 60}{13} = 60 \mu C$$

Charge on each capacitor is same

∴ they are in series.


9. A parallel plate capacitor with plate area A and plate separation d = 2 m has a capacitance of 4μF. The new capacitance of the system if half of the space between them is filled with a dielectric material of dielectric constant K = 3 (as shown in the figure) will be:





- Sol.** (A)  $2\mu\text{F}$  (B)  $32\mu\text{F}$  (C)  $6\mu\text{F}$  (D)  $8\mu\text{F}$

$$C_{\text{original}} = \frac{A\epsilon_0}{d}$$



$$C_1 = \frac{A\epsilon_0}{d/2} = \frac{2A\epsilon_0}{d} = C$$

$$C_2 = \frac{KA\epsilon_0}{d/2} = \frac{2KA\epsilon_0}{d} = \frac{6A\epsilon_0}{d} = 3C$$

$C_1$  &  $C_2$  are in series

$$C_{\text{new}} = \frac{C_1 C_2}{C_1 + C_2} = \frac{C \times 3C}{C + 3C} = \frac{3C}{4} = \frac{3}{4} \times \frac{2A\epsilon_0}{d} = \frac{3}{2} \times \frac{A\epsilon_0}{d}$$

$$C_{\text{new}} = \frac{3}{2} C_{\text{original}}$$

$$= \frac{3}{2} \times 4 = 6\mu\text{F}$$

- 10.** Sixty four conducting drops each of radius  $0.02\text{ m}$  and each carrying a charge of  $5\mu\text{C}$  are combined to form a bigger drop. The ratio of surface density of bigger drop to the smaller drop will be:

- Sol.** (A)  $1 : 4$  (B)  $4 : 1$  (C)  $1 : 8$  (D)  $8 : 1$

Let  $R$  = radius of combined drop

$r$  = radius of smaller drop

Volume will remain same

$$\frac{4}{3}\pi R^3 = 64 \times \frac{4}{3}\pi r^3$$

$$R = 4r; Q = 64q;$$

$q$  : charge of smaller drop


$Q$  : Charge of combined drop

$$\frac{\sigma_{\text{bigger}}}{\sigma_{\text{smaller}}} = \frac{\frac{Q}{4\pi R^2}}{\frac{q}{4\pi r^2}} = \frac{Q}{q} \cdot \frac{r^2}{R^2}$$

$$64 \frac{r^2}{16r^2} = 4 \Rightarrow \frac{\sigma_{\text{bigger}}}{\sigma_{\text{smaller}}} = \frac{4}{1}$$




11. The equivalent resistance between points A and B in the given network is:



- (A)  $65\Omega$                       (B)  $20\Omega$                       (C)  $5\Omega$                       (D)  $2\Omega$

Sol. (C)



$$R_{AB} = 5\Omega$$

12. A bar magnet having a magnetic moment of  $2.0 \times 10^5 \text{ JT}^{-1}$ , is placed along the direction of uniform magnetic field of magnitude  $B = 14 \times 10^{-5} \text{ T}$ . The work done in rotating the magnet slowly through  $60^\circ$  from the direction of field is:


- (A) 14J                      (B) 8.4J                      (C) 4J                      (D) 1.4J





**Sol. (A)**  
 Work done =  $MB (\cos \theta_1 - \cos \theta_2)$   
 $\theta_1 = 0^\circ, \theta_2 = 60^\circ$   
 $= 2 \times 10^5 \times 14 \times 10^{-5} (1 - 1/2)$   
 $= 14\text{J}$

**13.** Two coils of self inductance  $L_1$  and  $L_2$  are connected in series combination having mutual inductance of the coils as  $M$ . The equivalent self inductance of the combination will be:



- (A)  $\frac{1}{L_1} + \frac{1}{L_2} + \frac{1}{M}$       (B)  $L_1 + L_2 + M$       (C)  $L_1 + L_2 + 2M$       (D)  $L_1 + L_2 - 2M$

**Sol. (D)**  
 Current on both the inductor is in opposite direction.  
 Hence :  
 $L_{eq} = L_1 + L_2 - 2M$

**14.** A metallic conductor of length 1m rotates in a vertical plane parallel to east-west direction about one of its end with angular velocity  $5 \text{ rad s}^{-1}$ . If the horizontal component of earth's magnetic field is  $0.2 \times 10^{-4}\text{T}$ , then emf induced between the two ends of the conductor is:

- (A)  $5\mu\text{V}$       (B)  $50\mu\text{V}$       (C)  $5\text{mV}$       (D)  $50\text{mV}$

**Sol. (B)**  
 emf induced between the two ends =  $\frac{B_H \omega \ell^2}{2}$   
 $\frac{0.2 \times 10^{-4} \times 5 \times 1}{2} = 0.5 \times 10^{-4} = 50 \times 10^{-6} \text{ V} = 50\mu\text{V}$

**15.** Which is the correct ascending order of wavelengths?

- (A)  $\lambda_{\text{visible}} < \lambda_{\text{X-ray}} < \lambda_{\text{gamma-ray}} < \lambda_{\text{microwave}}$       (B)  $\lambda_{\text{gamma-ray}} < \lambda_{\text{X-ray}} < \lambda_{\text{visible}} < \lambda_{\text{microwave}}$   
 (C)  $\lambda_{\text{X-ray}} < \lambda_{\text{gamma-ray}} < \lambda_{\text{visible}} < \lambda_{\text{microwave}}$       (D)  $\lambda_{\text{microwave}} < \lambda_{\text{visible}} < \lambda_{\text{gamma-ray}} < \lambda_{\text{X-ray}}$

**Sol. (B)**  
 From electromagnetic wave spectrum.  
 $\lambda$  increases  $\rightarrow$

$\gamma$ - ray	x - rays	ultra violet	visible	infrared	microwave	Radio wave
----------------	----------	--------------	---------	----------	-----------	------------

$$\lambda_{\text{gamma-ray}} < \lambda_{\text{X-ray}} < \lambda_{\text{visible}} < \lambda_{\text{microwave}}$$

**16.** For a specific wavelength 670 nm of light coming from a galaxy moving with velocity  $v$ , the observed wavelength is 670.7 nm.

The value of  $v$  is :

- (A)  $3 \times 10^8 \text{ ms}^{-1}$       (B)  $3 \times 10^{10} \text{ ms}^{-1}$       (C)  $3.13 \times 10^5 \text{ ms}^{-1}$       (D)  $4.48 \times 10^5 \text{ ms}^{-1}$

**Sol. (C)**  
 $\lambda_{\text{emitted}} = 670 \text{ nm}$   
 $\lambda_{\text{obs}} = 670.7 \text{ nm}$   
 $v = ?$   
 $c = 3 \times 10^8 \text{ m/s}$   
 If  $v \ll c$   
 $\frac{\lambda_{\text{obs}} - \lambda_{\text{emitted}}}{\lambda_{\text{emitted}}} = \frac{v}{c} \Rightarrow \frac{670.7 - 670}{670} = \frac{v}{3 \times 10^8}$   
 $v = 3.13 \times 10^5 \text{ m/s}$



17. A metal surface is illuminated by a radiation of wavelength  $4500\text{\AA}$ . The ejected photo-electron enters a constant magnetic field of  $2\text{mT}$  making an angle of  $90^\circ$  with the magnetic field. If it starts revolving in a circular path of radius  $2\text{mm}$ , the work function of the metal is approximately:

(A)  $1.36\text{eV}$                       (B)  $1.69\text{eV}$                       (C)  $2.78\text{eV}$                       (D)  $2.23\text{eV}$

Sol. (A)

$$\lambda = 4500\text{\AA}$$

$$B = 2\text{mT}, R = 2\text{mm}$$

$$R = \frac{\sqrt{2Km}}{qB}$$

$$\frac{(qBR)^2}{2m} = K$$

$$\frac{(1.6 \times 10^{-19} \times 2 \times 10^{-3} \times 2 \times 10^{-3})^2}{2 \times 9.1 \times 10^{-31}} = K$$

$$\frac{(6.4)^2}{2 \times 9.1} \times \frac{10^{-50}}{10^{-31}} = K$$

$$K = 2.25 \times 10^{-19}\text{J}$$

$$= \frac{2.25 \times 10^{-19}}{1.6 \times 10^{-19}}\text{eV} = 1.40\text{eV}$$

$$E = \frac{12400}{4500} = 2.76\text{eV}$$

$$\phi = E - K = (2.76 - 1.40)\text{eV} = 1.36\text{eV}$$

18. A radioactive nucleus can decay by two different processes. Half-life for the first process is  $3.0$  hours while it is  $4.5$  hours for the second process. The effective half-life of the nucleus will be:

(A)  $3.75$  hours                      (B)  $0.56$  hours                      (C)  $0.26$  hours                      (D)  $1.80$  hours

Sol. (D)

$$\lambda_{\text{eq}} = \lambda_1 + \lambda_2$$

$$\frac{\ln 2}{(t_{1/2})_{\text{eq}}} = \frac{\ln 2}{(t_{1/2})_1} + \frac{\ln 2}{(t_{1/2})_2}$$

$$(t_{1/2})_{\text{eq}} = \frac{(t_{1/2})_1 \times (t_{1/2})_2}{(t_{1/2})_1 + (t_{1/2})_2}$$

$$= \frac{3 \times 4.5}{3 + 4.5} = \frac{3 \times 4.5}{7.5} = \frac{3 \times 3}{5} = 1.8\text{hr}$$

19. The positive feedback is required by an amplifier to act an oscillator. The feedback here means:

(A) External input is necessary to sustain ac signal in output.  
 (B) A portion of the output power is returned back to the input.  
 (C) Feedback can be achieved by LR network.  
 (D) The base-collector junction must be forward biased.

Sol. (B)

When the amplifier connects with positive feedback, it acts as the oscillator the feedback here is positive feedback which means some amount of voltage is given to the input.

20. A sinusoidal wave  $y(t) = 40\sin(10 \times 10^6 \pi t)$  is amplitude modulated by another sinusoidal wave  $x(t) = 20\sin(1000\pi t)$ . The amplitude of minimum frequency component of modulated signal is:

(A)  $0.5$                                       (B)  $0.25$                                       (C)  $20$                                       (D)  $10$



**Sol. (D)**

$$y(t) = 40\sin(10 \times 10^6 \pi t)$$

$$x(t) = 20\sin(1000 \pi t)$$

$$\Rightarrow \omega_c = 10^7 \pi$$

$$\omega_m = 10^3 \pi$$

$$A_c = 40$$

$$A_m = 20$$

$$\text{Equation of modulated wave} = (A_c + A_m \sin \omega_m t) \sin \omega_c t$$

$$= A_c \left( 1 + \frac{A_m}{A_c} \sin \omega_m t \right) \sin \omega_c t$$

$$A_c(1 + m \sin \omega_m t) \sin \omega_c t, \quad \mu = \frac{A_m}{A_c}$$

$$= A_c \sin \omega_c t + \frac{\mu A_c}{2} [\cos(\omega_c - \omega_m)t - \cos(\omega_c + \omega_m)t]$$

$$\text{Amplitude of minimum frequency} = \frac{\mu A_c}{2} = \frac{A_m}{A_c} \times \frac{A_c}{2} = \frac{A_m}{2} = 10$$

- 21.** A ball is projected vertically upward with an initial velocity of  $50 \text{ms}^{-1}$  at  $t = 0 \text{s}$ . At  $t = 2 \text{s}$ , another ball is projected vertically upward with same velocity. At  $t = \underline{\hspace{2cm}}$  s, second ball will meet the first ball ( $g = 10 \text{ms}^{-2}$ )

**Sol. (6)**

Let they meet at  $t = t$

So first ball gets  $t$  sec.

& 2<sup>nd</sup> gets  $(t - 2)$  sec. & they will meet at same height

$$h_1 = 50t - \frac{1}{2}gt^2$$

$$h_2 = 50(t - 2) - \frac{1}{2}g(t - 2)^2$$

$$h_1 = h_2$$

$$50t - \frac{1}{2}gt^2 = 50(t - 2) - \frac{1}{2}g(t - 2)^2$$

$$100 = \frac{1}{2}g[t^2 - (t - 2)^2]$$

$$100 = \frac{10}{2}[4t - 4]$$

$$5 = t - 1$$

$$t = 6 \text{sec.}$$

- 22.** A batsman hits back a ball of mass  $0.4 \text{kg}$  straight in the direction of the bowler without changing its initial speed of  $15 \text{ms}^{-1}$ . The impulse imparted to the ball is  $\underline{\hspace{2cm}}$  Ns.

**Sol. (12)**


Impulse = change in momentum

$$= m[v - (-v)] = 2mv$$


$$= 2 \times 0.4 \times 15 = 12 \text{ Ns}$$




23. A system to 10 balls each of mass 2kg are connected via massless and un-stretchable string. The system is allowed to slip over the edge of a smooth table as shown in the figure. Tension on the string between the 7<sup>th</sup> and 8<sup>th</sup> ball is \_\_\_\_\_ N when 6<sup>th</sup> ball just leaves the table.



Sol. (36)



$$a = \frac{6mg}{10m} = \frac{6g}{10} = \frac{3g}{5}$$



taking 8, 9, 10 together

$$T = 3ma$$

$$= 3m \times \frac{3g}{5}$$

$$= 36 \text{ N}$$

24. A geyser heats water flowing at a rate of 2.0 kg per minute from 30°C to 70°C. If geyser operates on a gas burner, the rate of combustion of fuel will be \_\_\_\_\_ g min<sup>-1</sup>. [Heat of combustion = 8 × 10<sup>3</sup> Jg<sup>-1</sup>]; Specific heat of water = 4.2 Jg<sup>-1</sup>°C<sup>-1</sup>]

Sol. (42)

$$m = 2000 \text{ gm/min}$$

$$\text{Heat required by water/min} = mS\Delta T$$

$$= (2000) \times 4.2 \times 40 \text{ J/min.}$$

$$= 336000 \text{ J/min.}$$

$$\left(\frac{dm}{dt}L\right) = 336000 \text{ J/min.}$$

$$\text{The rate of combustion} = \frac{dm}{dt} = \frac{336000}{8 \times 10^3} \text{ g/min.}$$

$$= 42 \text{ gm/min.}$$



25. A heat engine operates with the cold reservoir at temperature 324K. The minimum temperature of the hot reservoir, if the heat engine takes 300 J heat from the hot reservoir and delivers 180 J heat to the cold reservoir per cycle, is \_\_\_\_\_ K.


Sol. (540)

$$T_c = 324K$$

$$T_H = ?$$

$$Q_H = 300J$$

$$Q_L = 180 J$$



$$1 - \frac{Q_L}{Q_H} = 1 - \frac{T_L}{T_H}$$

$$\frac{Q_L}{Q_H} = \frac{T_L}{T_H}$$

$$T_H = \frac{Q_H}{Q_L} \times T_L = \frac{300}{180} \times 324 = 540K$$

26. A set of 20 tuning forks is arranged in a series of increasing frequencies. If each fork gives 4 beats with respect to the preceding fork and the frequency of the last fork is twice the frequency of the first, then the frequency of last fork is \_\_\_\_\_ Hz.

Sol. (152)

$$f_1 = f$$

$$f_2 = f + 4$$

$$f_3 = f + 2 \times 4$$

$$f_4 = f + 3 \times 4$$

$$f_{20} = f + 19 \times 4$$

$$f + (19 \times 4) = 2 \times f$$

$$f = 76Hz.$$

$$\text{Frequency of last tuning forks} = 2f$$

$$= 152 \text{ Hz}$$






27. Two 10 cm long, straight wires, each carrying a current of 5A are kept parallel to each other. If each wire experienced a force of  $10^{-5}$  N, then separation between the wires is \_\_\_\_\_ cm.

Sol. (5)

It should be mentioned, 10 cm wire is part of long wire.

Force experienced by unit length of wire

$$= \frac{\mu_0 I_1 I_2}{2\pi d}, I_1 = I_2 = 5A$$



Force experienced by wires of length 10cm

$$= \frac{\mu_0 I_1 I_2}{2\pi d} \times 10 \times 10^{-2}$$

$$10^{-5} = \frac{2 \times 10^{-7} \times 5 \times 5}{d} \times 10 \times 10^{-2}$$

$$d = 50 \times 10^{-3} \text{ m}$$


$$d = 50 \times 10^{-1} \text{ cm} = 5 \text{ cm}$$

28. A small bulb is placed at the bottom of a tank containing water to a depth of  $\sqrt{7}$  m. The refractive index of water is  $\frac{4}{3}$ . The area of the surface of water through which light from the bulb can emerge out is  $x\pi\text{m}^2$ .

The value of x is \_\_\_\_\_.

Sol. (9)

C : Critical angle



$$\tan C = \frac{r}{h}$$

$$r = h \tan C$$

$$\sin C = \frac{1}{\mu} = \frac{3}{4}$$

$$\tan C = \frac{3}{\sqrt{7}}$$

$$r = \sqrt{7} \times \frac{3}{\sqrt{7}} = 3$$

$$\text{Area of surface} = \pi r^2 = 9\pi\text{m}^2$$

29. A travelling microscope is used to determine the refractive index of a glass slab. If 40 divisions are there in 1cm on main scale and 50 Vernier scale divisions are equal to 49 main scale divisions, then least count of the travelling microscope is \_\_\_\_\_  $\times 10^{-6}$ m.



Sol. (5)

$$50\text{VSD} = 49\text{MSD}$$

$$1\text{VSD} = \frac{49}{50}\text{MSD}$$

$$\text{Least count} = 1\text{MSD} - 1\text{VSD}$$

$$= \left(1 - \frac{49}{50}\right)\text{MSD} = \frac{1}{50}\text{MSD}$$

$$1\text{MSD} = \frac{1}{40}\text{cm}$$

$$\text{Least count} = \frac{1}{50 \times 40}\text{cm}$$

$$\frac{1}{2000}\text{cm} = \frac{1}{2} \times 10^{-5}\text{m}$$

$$0.5 \times 10^{-5}\text{m}$$

$$5 \times 10^{-6}\text{m}$$

30. The stopping potential for photoelectrons emitted from a surface illuminated by light of wavelength  $6630\text{\AA}$  is  $0.42\text{ V}$ . If the threshold frequency is  $x \times 10^{13}\text{ /s}$ , where  $x$  is \_\_\_\_\_ (nearest integer) (Given, speed light  $3 \times 10^8\text{ m/s}$ , Planck's constant =  $6.63 \times 10^{-34}\text{ Js}$ )

Sol. (35)

$$\text{Stopping potential } V_0 = 0.42\text{V}$$

$$\lambda = 6630\text{\AA}$$

$$E = \phi + eV_0$$

$E$  : energy of incident photon

$V_0$  : Stopping potential

$$\phi = E - eV_0$$

$$E = \frac{12400}{6630}\text{eV} = 1.87\text{eV}$$

$$\phi = (1.87 - 0.42) = 1.45\text{eV}$$

$\phi = h\nu_0$  :  $\nu_0$  : threshold frequency


$$1.45 \times 1.6 \times 10^{-19} = 6.63 \times 10^{-34} \times \nu_0$$

$$\nu_0 = 0.35 \times 10^{15}$$

$$= 35 \times 10^{13}\text{ sec}^{-1} \Rightarrow x = 35$$



Perfect mix of  
**CLASSROOM** Program aided  
with technology for sure **SUCCESS.**



Continuing the legacy  
for the **last 15 years**



**MOTION LEARNING APP**

Get 7 days **FREE** trial & experience Kota Learning



Admission **OPEN**

**Umeed**  
**RANK KI HO**  
**ya Selection ki,**  
**Jeet Nischit hai**



Based on **JEE MAIN** June'22 Result

%tile Basis	Scholarship**
>99.9	90%
>99.5	75%
>99	50%
>98	40%
>97	30%
>95	20%

\* Scholarship calculation will be considered once the result is declared.  
\*\* Minimum amount is needed to be deposit at the time of admission.

सफलता की शुरुआत, सिर्फ मोशन के साथ...

Appear in instant

**MOTION<sup>®</sup> OPEN**  
**SCHOLARSHIP TEST**

Avail upto **100% SCHOLARSHIP** on  
**JEE, NEET & Foundation** (Class 6th-12th Pass) Courses

**MOTION<sup>®</sup>**  
JEE | NEET | NTSE | BOARDS | OLYMPIADS

 **1800 212 1799**